



Original article

<https://doi.org/10.18019/1028-4427-2023-29-5-552-556>**Combination of flexible intramedullary nailing and Ilizarov frame for salvage of femur and humerus nonunion in a girl with osteogenesis imperfecta**E.R. Mingazov¹, P. Foster², A.V. Popkov¹, D.A. Popkov^{1✉}¹ Ilizarov National Medical Research Centre for Traumatology and Orthopedics, Kurgan, Russian Federation² Leeds Teaching Hospitals NHS Trust, Leeds, UK**Corresponding author:** Dmitry A. Popkov, dpopkov@mail.ru**Abstract**

Background Fixation of pathological long bones with telescopic intramedullary rods is well known to be a technically challenging procedure even in specialist centres, with a high complication rate due to rod migration, hardware failure, nonunion or malunion. However there is very little guidance in the literature regarding salvage treatment options when failure occurs. **Aim** We demonstrate a surgical technique that can be used for salvage treatment of both femoral and humeral complex nonunions following Fassier-Duval (FD) rodding in a child with osteogenesis imperfecta (OI). **Case description** A 13 year-old girl with OI type VIII presented sequentially with nonunion and deformity of the femur then the humerus following previous FD rods in those segments. The femur was also complicated with metallosis between the steel rod and an overlying titanium plate. Both segments were treated with pseudarthrosis debridement, removal of metalwork and stabilisation with hydroxyapatite (HA)-coated flexible intramedullary nails, with temporary Ilizarov frame to provide enough longitudinal and rotational stability to allow immediate weight-bearing. The femur Ilizarov frame was removed after 64 days, and the femur remained straight and fully healed at 2.5 years. The frame time for the humerus was 40 days, complete union was achieved and upper limb function restored and maintained at 9 months. **Discussion** The transphyseal telescopic rod is the traditional implant of choice in terms of treating fractures and stabilising osteotomies for deformity in OI. However, it does not provide enough torsional or longitudinal stability by itself to allow early weight-bearing which is detrimental to bone healing in this vulnerable patient group. The incidence of delayed union or nonunion at osteotomy site in telescopic rod application is not negligible: up to 14.5-51.5 %. Although the technique we have shown in this case may not be applied to all complex OI patients, we believe that the combination of flexible intramedullary nails and Ilizarov frame provides a favourable environment for bone healing in complex or revision cases. As a secondary learning point the initial revision surgery to the left femur demonstrated the perils of using a steel rod and a titanium plate in a biologically active environment which in this case lead to metallosis and lysis. **Conclusion** We found the technique of HA-coated flexible intramedullary nails combined with the Ilizarov frame effective in the salvage of failed telescopic rods in both femur and humerus and feel this technique can be used as a salvage option in similar cases worldwide. This case also demonstrates the perils of using different metals in combined internal fixation.

Keywords: osteogenesis imperfecta type VIII, flexible intramedullary nailing, external fixation, telescopic rodding**For citation:** Mingazov E.R., Foster P., Popkov A.V., Popkov D.A. Combination of flexible intramedullary nailing and Ilizarov frame for salvage of femur and humerus nonunion in a girl with osteogenesis imperfecta. *Genij Ortopedii*. 2023;29(5):552-556. doi: 10.18019/1028-4427-2023-29-5-552-556**Научная статья**

УДК [616.71-007.235-001.59]-089.227.84

Комбинированное применение эластичного интрамедуллярного армирования и аппарата Илизарова в лечении ложного сустава бедра и плеча у девочки с несовершенным остеогенезомЭ.Р. Мингазов¹, Р. Foster², А.В. Попков¹, Д.А. Попков^{1✉}¹ Национальный медицинский исследовательский центр травматологии и ортопедии имени академика Г.А. Илизарова, Курган, Россия² Leeds Teaching Hospitals NHS Trust, Leeds, UK**Автор, ответственный за переписку:** Дмитрий Арнольдович Попков, dpopkov@mail.ru**Аннотация**

Введение. Остеосинтез телескопическими стержнями длинных трубчатых костей у детей при патологических состояниях костной ткани, включая несовершенный остеогенез (НО) – достаточно сложное хирургическое вмешательство со значимой частотой осложнений, обусловленных ротационной и продольной нестабильностью остеосинтеза, и, как следствие, с формированием несращений, ложных суставов или вторичных деформаций. **Цель.** Демонстрация хирургического лечения псевдоартрозов плечевой и бедренной костей у ребенка с НО, которому ранее была проведена коррекция деформаций остеосинтезом телескопическим стержнем Fassier-Duval. **Материалы и методы.** Девочка (13 лет) с VIII типом НО последовательно была пролечена по поводу несращений правого плеча и левого бедра, развившихся после ранее выполненных корригирующих остеотомий соответствующих сегментов и остеосинтеза телескопическими стержнями Fassier-Duval. Ложный сустав бедра также был осложнен металлозом вследствие попытки дополнительной стабилизации положения костных фрагментов короткой наконечной титановой пластиной. **Результаты.** Нами выполнены схожие операции: удаление предшествующего материала остеосинтеза, экономная резекция концов отломков, остеосинтез интрамедуллярными эластичными стержнями с остеиндуцирующим покрытием и редуцированным внешним остеосинтезом с целью обеспечения ротационной и продольной стабильности и возможности ранней нагрузки оперированных сегментов. Длительность внешней фиксации бедра составила 64 дня, костное сращение и коррекция деформации достигнуты. При контрольном осмотре через 2,5 года результат полностью сохраняется. Внешний остеосинтез плеча осуществлен в течение 40 дней, костное сращение достигнуто. При осмотре через 9 месяцев результат признан хорошим с полным функциональным восстановлением. **Обсуждение.** Телескопический трансфизарный остеосинтез признан методом выбора при оперативном лечении деформаций длинных трубчатых костей у детей с НО. Однако данный способ не обеспечивает достаточной ротационной и продольной стабильности костных фрагментов, что может являться источником вторичных осложнений. Частота несращений достигает 14,5-51,5 %. В сложных и ревизионных случаях деформаций и ложных суставов при НО мы видим преимущества комбинированного остеосинтеза (внутренняя и редуцированная внешняя фиксация) в создании благоприятных условий для костного сращения и ранней функциональной нагрузки. На основании данного случая также подчеркнем недопустимость одновременного применения в очаге остеосинтеза имплантов из разных металлов (стальной и титановый сплав). **Заключение.** Данный случай показывает эффективность применения комбинации эластичных интрамедуллярных стержней с остеиндуцирующим покрытием и внешней фиксации по Илизарову при лечении ложных суставов после ранее проведенного лечения у детей с НО.

Ключевые слова: несовершенный остеогенез тип VIII, эластичное интрамедуллярное армирование, внешняя фиксация, телескопический остеосинтез**Для цитирования:** Мингазов Э.Р., Foster P., Попков А.В., Попков Д.А. Комбинированное применение эластичного интрамедуллярного армирования и аппарата Илизарова в лечении ложного сустава бедра и плеча у девочки с несовершенным остеогенезом. *Гений ортопедии*. 2023;29(5):552-556. doi: 10.18019/1028-4427-2023-29-5-552-556. EDN: RNSLCK.

© Mingazov E.R., Foster P., Popkov A.V., Popkov D.A., 2023

BACKGROUND

Osteogenesis imperfecta (OI) is a phenotypically and genetically heterogeneous group of inherited bone dysplasias characterized by frequent fractures, bone deformities, low bone mineral density and osteopenia [1-4]. Although rare, each patient often requires multiple operations on several limbs throughout childhood and beyond, in order to maintain autonomy, self-care, physical activity and the ability to acquire and develop motor skills including independent ambulation.

The workhorse implant for the treating surgeon in cases of OI is the telescopic rod which strengthens and protects the full length of the long bones, where one component (male) slides inside the other (female) with growth. As well as allowing fractures and osteotomies to heal the rods prevent deformity and fracture recurrence [5, 6]. Models include the Fassier-Duval rod [7-9], the Sheffield rod [10], dual interlocking telescopic rod [11], and the titanium telescopic rod [12, 13].

Whatever the exact implant design telescopic rodding carries a steep learning curve, and even in the

most experienced hands has a high complication rate, partly related to the lack of rotational and longitudinal stability of the implant [6, 11, 14, 15]. The technique is known to be even more difficult and prone to these complications in the humerus, including secondary displacement, malunion, nonunion, hardware failure and migration over time and bone growth [16-19]. This means that revision surgery in these cases is not at all unusual, and is even more challenging than primary cases requiring individual solutions as well as classic techniques [9, 18, 20-22].

Aim This article presents a case report of operative treatment of a 13-year-old girl affected with osteogenesis imperfecta type VIII, who was consecutively operated on with a combined technique with hydroxyapatite-coated flexible intramedullary nailing (HA-FIN) and reduced external frame for salvage of femoral pseudarthrosis complicated with metallosis and pseudarthrosis of humerus after Fassier-Duval (FD) rod failure in both segments.

CASE DESCRIPTION

The initial treatment of this child with regards to the right femur was described in a published report [18]. Two years later the patient represented to us with a symptomatic nonunion in the other (left) femur, having been treated with primary FD rod elsewhere in 2013, and then attempted revision with exchange FD rod followed by additional titanium plate 4 months before presentation.

On review, the girl was unable to walk or stand, on a background of short stature with rhizomelia, white sclerae, myopia and partial hearing loss. Pain was localized to the midshaft of the left femur and knee, in the area of the mobile nonunion site. There was a leg length discrepancy of 1.5 cm and 25 degree loss of extension in the left knee. There was no evidence of infection including on blood tests. Clinical alignment of the right lower limb remained excellent without any deformity rebound or implant migration, with satisfactory X-rays. Views of left femur showed a clear nonunion with a 4 mm gap, with FD rod in place as well as lysis around the supplementary plate and screws.

Left femur surgery

The existing FD rod was removed as well as open debridement of the pseudarthrosis and overlying plate and screws. Extensive metallosis was noted around the soft tissues and periosteum around the plate. The canal was reamed to remove any bone fragments from the screw tracts and the previous apex anterior deformity corrected. Two 2.5 mm HA-coated titanium flexible intramedullary nails (Orthopediatrics nails modified by Metis Ltd, Tomsk, Russia) were inserted in anterograde fashion. The Ilizarov apparatus (Experimental Plant at the Russian Ilizarov Scientific Center for Restorative Traumatology and Orthopaedics, Kurgan, Russia) consisted of a short proximal arch with three half-pins and a distal ring with three wires. This was a static construct which did not require any postoperative adjustment or correction.

Standing and progressive weight-bearing commenced on day 3 and the patient was discharged home on day 8 with a physical therapy programme including early knee

movement. Postoperatively, no problems or obstacles were encountered such as pin-site infection or loosening. X-rays at 60 days showed union and the fixator was removed at day 64 under general anaesthetic. The lower limb was protected in a split cast for a period of two weeks and then weight-bearing was recommenced. Bisphosphonate therapy was resumed at 4 months. Independent standing and pain-free walking was restored and maintained at 2 years, with full union and alignment of the femur demonstrated on X-rays with remodelling and restitution of the intramedullary canal.

Right humerus presentation and surgery

Shortly after this episode there was a similar presentation of the right arm in the same child. The right humerus had been previously treated in another institution with FD rod. A painful and mobile non-union of the distal humerus was noted, with functional impairment. X-rays revealed an established nonunion as well as broken male component of the FD rod. There was no evidence of infection including on blood tests.

A proximal approach was used to remove both the female component of the FD rod as well as the proximal part of the broken male component. The nonunion site was debrided back to viable bone and the distal part of the male component removed. In similar principles to the femur surgery, the humerus was stabilised with two HA-coated flexible intramedullary nails and a reduced Ilizarov frame, consisting of 4 half-pins fixed over a short Ilizarov plate connected with threaded rods. This provided excellent bone contact, rotational stability and allowed free elbow movement. Neither casting nor bracing was required or any postoperative adjustments to the Ilizarov apparatus.

There were no problems or obstacles postoperatively. The humerus healed and the fixator was removed at 40 days, and the arm was left free to move. Bisphosphonate therapy was recommenced at 6 months. At 9 months there was pain-free restoration of upper limb function, as well as radiological union and remodelling of the canal.




Fig. 1. Preoperative radiographs: a – pseudarthrosis of the left femur and hardware fracture; b – fractured rod has been replaced with another Fassier-Duval rod, note interfragmentary diastasis (fleshs) as result of longitudinal instability; c – standing radiograph of lower limbs, augmented radiographs demonstrate loosening of the plate and bicortical screws, nonunion




Fig. 2. Treatment of left femur pseudarthrosis: a – intraoperative view: nonunion site extensive metallosis of multiple charred appearance inclusions on the intraoperative view, removed hardware; b – postoperative X-rays; c – day 64 after surgery, radiographs at frame removal; d – 2 years after frame removal from left femur




Fig. 3. Treatment of right humerus pseudarthrosis: a – preoperative radiographs, pseudarthrosis and broken Fassier-Duval rod; b – postoperative radiographs; c – bone union achieved, frame removed on day 40; d – 9 months after frame removal from right humerus



Fig. 4. Functional outcome

DISCUSSION

The transphyseal telescopic rod is the traditional implant of choice in terms of treating fractures and stabilising osteotomies for deformity in OI [5-8]. However, it does not provide enough torsional or longitudinal stability to allow early weight-bearing [6, 8, 10, 11, 16] which is detrimental to bone healing in this vulnerable patient group [23, 24] as well as the overall rehabilitation [22].

The incidence of delayed union or nonunion at osteotomy site in telescopic rod application is not negligible: up to 14.5 % [16]. Munns et al observed delayed bone union in 103 cases out of 200 (51.5 %) operated on with the Fassier-Duval rod [17]. Shin et al. reported 5 cases of persisting cortical gap for a series of 59 rods [11]. The case presented in our article does not match to reported data. Nonunion and rod fracture occurred in two segments of three within period of 3 and 5 years. The peer-reviewed literature reports that for a five-year period, the telescopic rods did not required surgery in 63 % for the femur and 64 % for tibia according to Cox et al [25]. Shin et al used Dual Interlocking telescopic rod (D-ITR) and reported 75 % survival rate for 5.3 years follow-up, which is the best result among all known telescopic rods [11]. On the other hand, the rate of revision surgery also depends on the severity of OI and it reaches 67.86 % in OI type III and 31.82 % in OI type IV [26]. We presume

that there could be features of bone union in patients with type VIII of OI (recessive inheritance pattern). There is no relevant literature for OI type VIII specifically.

Even in "classic" OI the literature suggests that corrective osteotomy of the humerus with FD rod carries a high complication rate [19, 20]. Grossman et al reported revision rate of 34.3 % within 35 months including unscheduled surgery for nonunion and malunion in 8.6 % of cases [19].

Although the technique we have shown in this case may not be applied to all complex OI patients (such as the very young) we believe that the combination of HA-coated flexible intramedullary nails and Ilizarov frame provides a favourable environment for bone healing in complex or revision cases [28, 29]. There is an enhanced mechanical stability (both particularly rotational and longitudinal) compared to telescopic rods alone, as well as a favourable osteoconductive environment.

As a secondary learning point, the initial revision surgery to the left femur demonstrated the perils of using a steel rod and a titanium plate in a biologically active environment which in this case lead to metallosis and lysis [30, 31]. The technique of combined telescopic rodding and plate fixation per se is supported by the literature, but the metals used should be similar, also locking unicortical screws rather than plates with bicortical conventional screws [15, 27].

CONCLUSION

We found a high rate of nonunion and pseudarthrosis associated with hardware fracture in a patient with rare type VIII of osteogenesis imperfecta initially treated with FD rods in each segment. A combination of pseudarthrosis site resection, elastic intramedullary nailing with osteoinductive coating and external frame

applied for a short period provides early function including full weight-bearing and enables bone union and further favorable bone remodeling in long-term follow-up. The surgeon should also be aware about the risks of use of dissimilar metals in combined osteosynthesis resulting in osteolysis and early loosening of implants.

Conflict of interest. Absent.

Funding Additional sources of funding were not attracted.

Ethical expertise Not required.

Informed consent Received.

REFERENCES

1. Sillence DO, Senn A, Danks DM. Genetic heterogeneity in osteogenesis imperfecta. *J Med Genet.* 1979;16(2):101-116. doi: 10.1136/jmg.16.2.101
2. Cheung MS, Glorieux FH. Osteogenesis Imperfecta: update on presentation and management. *Rev Endocr Metab Disord.* 2008;9(2):153-160. doi: 10.1007/s11154-008-9074-4
3. Van Dijk FS, Pals G, Van Rijn RR, et al. Classification of Osteogenesis Imperfecta revisited. *Eur J Med Genet.* 2010;53(1):1-5. doi: 10.1016/j.ejmg.2009.10.007
4. Glorieux FH. Osteogenesis imperfecta. *Best Pract Res Clin Rheumatol.* 2008;22(1):85-100. doi: 10.1016/j.berh.2007.12.012
5. Rodriguez Celin M, Kruger KM, Caudill A, et al. A Multicenter Study of Intramedullary Rodding in Osteogenesis Imperfecta. *JB JS Open Access.* 2020;5(3):e20.00031. doi: 10.2106/JBJS.OA.20.00031

6. Birke O, Davies N, Latimer M, et al. Experience with the Fassier-Duval telescopic rod: first 24 consecutive cases with a minimum of 1-year follow-up. *J Pediatr Orthop*. 2011;31(4):458-464. doi: 10.1097/BPO.0b013e31821bf50
7. Fassier F. Fassier-Duval Telescopic System: How I Do It? *J Pediatr Orthop*. 2017;37 Suppl 2:S48-S51. doi: 10.1097/BPO.0000000000001024
8. Suiko J, Oberc A. Advantages and Complications Following Fassier-Duval Intramedullary Rodding in Children. Pilot Study. *Ortop Traumatol Rehabil*. 2015;17(5):523-30. doi: 10.5604/15093492.1186830
9. Fassier FR. Osteogenesis Imperfecta-Who Needs Rodding Surgery? *Curr Osteoporos Rep*. 2021;19(3):264-270. doi: 10.1007/s11914-021-00665-z
10. Nicolaou N, Bowe JD, Wilkinson JM, et al. Use of the Sheffield telescopic intramedullary rod system for the management of osteogenesis imperfecta: clinical outcomes at an average follow-up of nineteen years. *J Bone Joint Surg Am*. 2011;93(21):1994-2000. doi: 10.2106/JBJS.J.01893
11. Shin CH, Lee DJ, Yoo WJ, et al. Dual Interlocking Telescopic Rod Provides Effective Tibial Stabilization in Children With Osteogenesis Imperfecta. *Clin Orthop Relat Res*. 2018;476(11):2238-2246. doi: 10.1097/CORR.0000000000000429
12. Popkov D, Dolganova T, Mingazov E, et al. Combined technique of titanium telescopic rods and external fixation in osteogenesis imperfecta patients: First 12 consecutive cases. *J Orthop*. 2020;22:316-325. doi: 10.1016/j.jor.2020.05.017
13. Mingazov ER, Gofman FF, Popkov AV, et al. First use experience with titanium telescopic rod in pediatric limb deformity correction in osteogenesis imperfecta. *Genij Ortopedii*. 2019;25(3):297-303. doi: 10.18019/1028-4427-2019-25-3-297-303
14. Sterian A, Balanescu R, Barbilian A, et al. Early telescopic rod osteosynthesis for Osteogenesis Imperfecta patients. *J Med Life*. 2015;8(4):544-547.
15. Cho TJ, Lee K, Oh CW, et al. Locking plate placement with unicortical screw fixation adjunctive to intramedullary rodding in long bones of patients with osteogenesis imperfecta. *J Bone Joint Surg Am*. 2015;97(9):733-737. doi: 10.2106/JBJS.N.01185
16. Azzam KA, Rush ET, Burke BR, et al. Mid-term Results of Femoral and Tibial Osteotomies and Fassier-Duval Nailing in Children With Osteogenesis Imperfecta. *J Pediatr Orthop*. 2018;38(6):331-336. doi: 10.1097/BPO.0000000000000824
17. Munns CF, Rauch F, Zeitlin L, et al. Delayed osteotomy but not fracture healing in pediatric osteogenesis imperfecta patients receiving pamidronate. *J Bone Miner Res*. 2004;19(11):1779-1786. doi: 10.1359/JBMR.040814
18. Popkov D. Use of flexible intramedullary nailing in combination with an external fixator for a postoperative defect and pseudarthrosis of femur in a girl with osteogenesis imperfecta type VIII: a case report. *Strategies Trauma Limb Reconstr*. 2018;13(3):191-197. doi: 10.1007/s11751-018-0320-3
19. Grossman LS, Price AL, Rush ET, et al. Initial Experience With Percutaneous IM Rodding of the Humeri in Children With Osteogenesis Imperfecta. *J Pediatr Orthop*. 2018;38(9):484-489. doi: 10.1097/BPO.0000000000000856
20. Hsiao MS, Mormino MA, Esposito PW, Burke BA. Distal humerus atrophic nonunion in a child with osteogenesis imperfecta. *J Pediatr Orthop*. 2013;33(7):725-729. doi: 10.1097/BPO.0b013e3182a32e69
21. Langlais T, Pannier S, De Tienda M, et al. 'In-Out-In' K-wires sliding in severe tibial deformities of osteogenesis imperfecta: a technical note. *J Pediatr Orthop B*. 2021;30(3):257-263. doi: 10.1097/BPB.0000000000000785
22. Fassier A. Telescopic rodding in children: Technical progression from Dubow-Bailey to Fassier-Duval™. *Orthop Traumatol Surg Res*. 2021;107(1S):102759. doi: 10.1016/j.otsr.2020.102759
23. Zeitlin L, Fassier F, Glorieux FH. Modern approach to children with osteogenesis imperfecta. *J Pediatr Orthop B*. 2003;12(2):77-87. doi: 10.1097/01.bpb.0000049567.52224.f
24. Esposito P, Plotkin H. Surgical treatment of osteogenesis imperfecta: current concepts. *Curr Opin Pediatr*. 2008;20(1):52-57. doi: 10.1097/MOP.0b013e3182f35f03
25. Cox I, Al Mouazzen L, Bleibleh S, et al. Combined two-centre experience of single-entry telescopic rods identifies characteristic modes of failure. *Bone Joint J*. 2020;102-B(8):1048-1055. doi: 10.1302/0301-620X.102B8.BJJ-2020-0131.R1
26. Rosemberg DL, Goiano EO, Akkari M, Santili C. Effects of a telescopic intramedullary rod for treating patients with osteogenesis imperfecta of the femur. *J Child Orthop*. 2018;12(1):97-103. doi: 10.1302/1863-2548.12.170009
27. Franzone JM, Kruse RW. Intramedullary nailing with supplemental plate and screw fixation of long bones of patients with osteogenesis imperfecta: operative technique and preliminary results. *J Pediatr Orthop B*. 2018;27(4):344-349. doi: 10.1097/BPB.0000000000000405
28. Boutaud B, Laville JM. Elastic sliding central medullary nailing with osteogenesis imperfecta. Fourteen cases at eight years follow-up. *Rev Chir Orthop Reparatrice Appar Mot*. 2004;90(4):304-311. (In French) doi: 10.1016/s0035-1040(04)70125-7
29. Mingazov ER, Popkov AV, Kononovich NA, et al. Results of using transphyseal elastic intramedullary nailing in patients with severe types of osteogenesis imperfecta. *Genij Ortopedii*. 2016;22(4):6-16.
30. Eliaz N. Corrosion of Metallic Biomaterials: A Review. *Materials* (Basel). 2019;12(3):407. doi: 10.3390/ma12030407
31. Zartman KC, Berlet GC, Hyer CF, Woodard JR. Combining dissimilar metals in orthopaedic implants: revisited. *Foot Ankle Spec*. 2011;4(5):318-323. doi: 10.1177/1938640011422597

The article was submitted 16.06.2023; approved after reviewing 26.06.2023; accepted for publication 25.08.2023.

Статья поступила в редакцию 16.06.2023; одобрена после рецензирования 26.06.2023; принята к публикации 25.08.2023.

Information about the authors:

1. Eduard R. Mingazov – Candidate of Medical Sciences, orthopedic surgeon, ed.dobryak@gmail.com, Scopus: 56897223100;
2. Patrick Foster – MD, PhD, pediatric orthopedic surgeon, patrick.foster@nhs.net, Scopus: 7201517406;
3. Arnold V. Popkov – Doctor of Medical Sciences, Professor, orthopedic surgeon, apopkov.46@mail.ru, https://orcid.org/0000-0001-5791-1989;
4. Dmitry A. Popkov – Doctor of Medical Sciences, Professor, orthopedic surgeon, dpopkov@mail.ru; https://orcid.org/0000-0002-8996-867X.

Информация об авторах:

1. Эдуард Рифович Мингазов – кандидат медицинских наук, врач-ортопед, ed.dobryak@gmail.com, Scopus: 56897223100;
2. Патрик Фостер – доктор медицинских наук, детский хирург-ортопед, patrick.foster@nhs.net, Scopus: 7201517406;
3. Арнольд Васильевич Попков – доктор медицинских наук, профессор, хирург-ортопед, apopkov.46@mail.ru, https://orcid.org/0000-0001-5791-1989;
4. Дмитрий Арнольдович Попков – доктор медицинских наук, профессор, хирург-ортопед, dpopkov@mail.ru, https://orcid.org/0000-0002-8996-867X.

Contribution of the authors:

Mingazov E.R. – Development, research activities; Application of formal methods for the analysis or synthesis of research data.
 Popkov A.V. – Responsibility for managing and coordinating the planning and conduct of research activities
 Foster P. – Preparation and writing of the initial draft (draft) of the work.
 Popkov D.A. – Gathering data/evidence, Preparing and writing an initial draft of the paper, Applying formal methods to synthesize study data.
 Conducting surgical treatment.

Вклад авторов:

Мингазов Э.Р. – разработка, проведение исследовательской деятельности; применение формальных методов для анализа или синтеза данных исследования.
 Попков А.В. – ответственность за управление и координацию планирования и проведения исследовательской деятельности.
 Foster P. – подготовка и написание первоначального проекта (черновика) работы.
 Попков Д.А. – сбор данных/доказательств, подготовка и написание первоначального проекта работы, применение формальных методов для синтеза данных исследования. Проведение оперативного лечения.